



Transformation of *Rhodiola rosea* with rol-genes from *Agrobacterium rhizogenes* to enhance the production of bioactive compounds and analyses of key genes in their pathways

Stofberg, Jimmi; Antypa, Natalia-Meropi; Müller, Renate; Lütken, Henrik Vlk

Publication date:
2015

Document version
Early version, also known as pre-print

Citation for published version (APA):
Stofberg, J., Antypa, N-M., Müller, R., & Lütken, H. V. (2015). *Transformation of Rhodiola rosea with rol-genes from Agrobacterium rhizogenes to enhance the production of bioactive compounds and analyses of key genes in their pathways*. Poster session presented at International Phytocosmetic Congress, Bogota, Colombia.

Transformation of *Rhodiola rosea* with *rol*-genes from *Agrobacterium rhizogenes*

– a platform to enhance production of the secondary metabolites salidroside and rosavin



Jimmi Stofberg, Natalia-Meropi Antypa, Renate Müller, Henrik Lütken

University of Copenhagen, Faculty of Science, Department of Plant and Environmental Sciences, Crop Sciences Section
Højbakkegård Allé 9-13, DK-2630 Taastrup, Denmark
tvr996@alumni.ku.dk

Key result

~60 % of the transformed explants developed hairy roots

Introduction

Rhodiola rosea L. (Fig. 1a) also known as roseroot is an arctic alpine plant distributed on the Northern hemisphere¹ (Fig. 1b). The plant contains among others the secondary metabolites rosavinoids and salidroside² (Fig. 2). These composites are active compounds that have been used for centuries to alleviate depression, stimulate the memory and as anti-fatigue and anti-inflammatory agents.

Transformation with *root oncogenic loci* (*rol*) genes from *Agrobacterium rhizogenes* leads to development of hairy roots at the infection site³. Transformed hairy roots have, for several plant species, been shown to contain higher contents of secondary metabolites compared to wild type⁴.

Objectives

- To obtain hairy roots of *R. rosea* containing *rol*-genes for future sustainable production of valuable bioactive compounds in bioreactors.
- To enhance the level of bioactive compounds *in planta*.
- To regenerate plants from transformed roots.




Figure 1. *Rhodiola rosea*. a) plant habitus b) geographical distribution of *R. rosea*




Figure 2 Structures of important secondary metabolites in *R. rosea*




Figure 3. Root formation following transformation. *R. rosea* was subjected to *A. rhizogenes* (transformed, n=74) or MYA (control, n =71).





Figure 4 Hairy root development. *R. rosea* was subjected to *A. rhizogenes* (transformed, n=74) or MYA (control, n =71)



Experimental design. The structure of the workflow behind the experiment for cultivation and regeneration of the putatively transformed hairy roots.

References:

- 1) Ramawat, K.G., 2008. Bioactive molecules and medicinal plants. In *Bioactive molecules and medicinal plants*. pp. 298–316.
- 2) Den virtuelle flora, 1997. Den virtuelle flora., <http://linnaeus.nrm.se/flora/di/crassula/rhodi/wel>.
- 3) Kiselev, K. V et al., 2007. The *rolB* gene-induced overproduction of resveratrol in *Vitis amurensis* transformed cells. *Journal of biotechnology*, 128(3), pp.681–92.
- 4) Britton, M.T. et al., 2013. Chapter 14 THE ONCOGENES OF AGROBACTERIUM TUMEFACIENS AND AGROBACTERIUM RHIZOGENES. *Springer*, pp.523–563.

Preliminary results

- Approximately 60 % of the transformed explants developed hairy roots (Fig. 3).
- Collectively the transformed explants developed hairy roots faster than the controls (Fig. 4).